DARI - KOOL THE VALUE LEADER 7 DARLKA

CHECK THESE OUTSTANDING FEATURES . . .

ON THE NEW NEW NEW DARI-KOOL J

Dari-Kool Direct Expansion Coolers with New Style Washing Unit

Refrigeration Systems

Heavy Duty Agitators

Built-In Washing System

Tube and Plate Pre-Coolers

Temperature Recorder with **New Alarm Circuit or Light**

Therma•Stor Heat Recovery System





ALL NEW DARI-KOOL-<u>The value</u> LEADER IN COOLING SYSTEMS

Dairymen.

Call 717-626-1151

For Details on These

Features On The

Now is the time

[Continued from Page 10]

continue for the next several weeks. Look for evidence of feeding activity on shade trees, oaks in particular as they are a favorite host tree. The feeding activity normally starts long before the average person is aware of their presence. They will start at the top of the tree and work down most of the feeding taking place at night.

Control measures will vary. A few shade trees can be fairly well protected by using burlap aprons around tree trunks. As the caterpillars crawl down the tree in the early morning, they will seek shelter under the burlap and can be killed. This needs daily attention.

Trees can also be protected with sprays. Since many trees are quite tall, this is a job that may be best done by a professional. Insecticides registered for use on Gypsy moth are Sevin and Dylox applied according to lable instructions.

Wooded areas are most succeptable but the high winds of last week will move them into new areas.

XXX To Check Soil Temperature For Soybean Planting

Soybeans should be planted in warm, moist soil. A thermometer is an ideal instrument in helping you decide when is the best planting date. Special bayonet type thermometers work best, however any wall type mercury thermometer will do.

The ideal soil temperature will be about 62 degrees F. Soil temperatures should be checked on at least two consecutive days. The reading should be taken at

about 8 a.m. when soil temperature has stabilized. The thermometer bulb should be inserted to a twoinch depth.

Full season soybeans will benefit most from early planting. This will allow a longer vegetative growth period before the start of the flowering period. Flowering is triggered by the shortened day length after June 21. This means maximum growth before June 21 will mean higher soybean yields.

XXX To Be Alert For Ticks

The tick season is here. These pests may be on almost any dog or person that walk through uncultivated fields or woody areas from May to September.

Ticks await their victims on low-growing shrubs and on tall grass. They attach to, and feed on the blood of dogs, humans as well as many other animals.

There are two common kinds of ticks, the American dog tick and the brown dog tick. Both ticks are brown, but the American dog tick has a mottled white shield on its back. The brown dog tick is a household pest and cannot survive our winters outdoors.

To control the American dog tick, first clean the area of tall grass and weeds on which the ticks await their victims. Also treat bushes 20 to 30 feet on either side of paths with either Sevin, Diazinon or Lindome. The dog should also be treated with a per cent Sevin dust at weekly intervals or as needed for control of either the American Dog or Brown Dog Tick.

Know Where the Activities Will Be? Read the Farm Women Calendar.

